

CLASS #6

**INDEPENDENT EVENTS AND
CONDITIONAL
PROBABILITIES**

Textbook's Sections 2.4 and 2.5
Probability, 1016-351

Rochester Institute of Technology

A Rule of Probability

$$P(A \text{ and } B) = P(A)P(B) \quad [\text{page 77}]$$

only for A and B independent events.

This is called the multiplication rule.

example: P(rain in London tomorrow
and Buffalo Bills will win the next
Super Bowl)

Example

Given that we know that $P(\text{rain in London tomorrow}) = 0.7$ and $P(\text{Buffalo Bills will win the next Super Bowl}) = 0.03$,

$P(\text{rain in London tomorrow}$
and $\text{Buffalo Bills will win the next}$
 $\text{Super Bowl})$


$$= (0.7)(0.03) = 0.021$$

the probability that both will occur

Independent events are events such that knowing about the occurrence or non-occurrence of one of them tells you nothing about the probability that the other event will occur.

A different type of example: Are
“getting an ace” and “getting a
spade” independent events
when randomly drawing a
single card from a standard
deck of cards?

YES _____ , NO _____



A Previous Example Revisited

We are given the three probabilities below

$$P(A) = P(\text{engineering major}) = 0.071$$

$$P(B) = P(\text{female}) = 0.320$$

$$P(A \text{ and } B) = P(\text{engineering major and female}) = 0.005$$

$$P(A \text{ or } B) = P(A) + P(B) - P(A \text{ and } B)$$

Example continued:

$$P(A \text{ or } B)$$

$$= P(\text{engineering major or female})$$

$$= 0.071 + 0.320 - 0.005 = 0.386$$

Are being a female and being an engineering major independent?

No.

$$P(\text{female})P(\text{engineer}) = (0.320)(0.071) \\ = 0.02272$$

$$P(\text{being both}) = 0.005$$

These would be the same values if they were independent events.

Conditional Probabilities

Example on the board:

Sample from a collection of coins containing 60 showing Phillip the Good and 40 showing Charles the Bold.

When selecting two coins, place the first coin aside before selecting the second coin. This is called sampling without replacement.

The following is an example from another textbook:

In a sample of 150 residents, each person was asked if he or she favored the concept of having a single countywide police agency. The county is composed of one large city and many suburban townships. The residence (city or outside the city) and the responses of the residents are summarized in Table 4.3. If one of these residents was to be selected at random, what is the probability that the person will (a) favor the concept? (b) favor the concept if the person selected is a city resident? (c) favor the concept if the person selected is a resident from outside the city? (d) Are the events F (favor the concept) and C (reside in city) independent?

Residence	Opinion		Total
	Favor (F)	Oppose (\bar{F})	
In city (C)	80	40	120
Outside of city (\bar{C})	20	10	30
<i>Total</i>	100	50	150

Note that this textbook uses \bar{F} for “not F.”

The entries in the table could be written as probabilities:

		Opinion		
		F	not F	
Residence	City	80/150	40/150	120/150
	Not City	20/150	10/150	30/150
		100/150	50/150	1

(a) $P(\text{Favor}) = 100/150$

(b) $P(\text{Favor} \text{ given that City}) = 80/120$

(c) $P(\text{Favor given that Outside the City})$
 $= 20/30$

(d) Are Favoring and being in the City independent? Answer: **Yes**,
since $P(F) P(C) = (100/150) (120/150)$
 $= 8/15 = 0.5333\dots$
and $P(F \text{ and } C) = 80/150 = 8/15$
 $= 0.5333\dots$

for Part (d), independence can be shown at least two other ways:

- (1) $P(F) = 100/150 = 2/3$ and $P(F \mid C) = 80/120 = 2/3$ from Parts (a) and (b), so $P(F) = P(F \mid C)$. That is, the probability of F occurring is the same as the probability of F occurring if C has occurred.
- (2) The rows of the table are proportional. That is 80 is to 40 as 20 is to 10.

Always (see page 69):

$$P(A \text{ and } B) = P(A) P(B | A)$$

Independent events (see page 77):

$$P(A \text{ and } B) = P(A) P(B)$$

A criteria for independence is

$$P(B | A) = P(B)$$

Some words that have been used

conditional probability

row

column

matrix notation

contingency table

cross classify

marginal

marginal probability

marginal sum

more →

Some words that have been used
continued

sample size

two way table

bivariate distribution

variables (Opinion and Residence)

r by c table (This example is 2 by 2.)

Some formulas that have been used

For any events E and F,

$$P(F \mid E) = P(E \text{ and } F) / P(E)$$

$$P(E \text{ and } F) = P(E) P(F \mid E)$$

For independent events E and F,


$$P(E \text{ and } F) = P(E) P(F)$$

$$P(F \mid E) = P(F)$$

Consider events E and F that have positive probability

If E and F are mutually exclusive, can they be independent?


Yes (always) ____ No (never) ____ Maybe ____



Consider events E and F that have positive probability

If E and F are independent, can they be mutually exclusive?

Yes (always) ____ No (never) ____ Maybe ____





LAST SLIDE FOR CLASS #6