

1016-345-01

Probability and Statistics for Engineers

Problem Set 8

Assigned 2012 October 30
Due 2012 November 6

Show your work on all problems! If you use a computer to assist with numerical computations, turn in your source code as well.

1 Devore Chapter 7, Problem 4

2 Devore Chapter 7, Problem 22

Note that problem 7.22 is different in the seventh and eighth editions of Devore. Be sure to do the problem from the eighth edition.

3 Devore Chapter 7, Problem 34

4 Devore Chapter 7, Problem 44

5 Computational Exercise (Extra Credit)

Suppose that a proportion p of the electorate prefers candidate A to candidate B, and a poll is conducted which selects a truly random sample of $n = 1000$ voters. The polling firm wishes to quote a 95% CL (i.e., $\alpha = .05$) interval on their estimate of p . If they find $n\hat{p}$ voters expressing a preference for candidate A, the lower and upper ends p_- and p_+ of the CI are given by

$$p_{\pm} = \frac{\hat{p} + z_{\alpha/2}^2/(2n)}{1 + z_{\alpha/2}^2/n} \pm \frac{z_{\alpha/2} \sqrt{\hat{p}(1 - \hat{p})/n + z_{\alpha/2}^2/(4n^2)}}{1 + z_{\alpha/2}^2/n}$$

- Plot p_- and p_+ versus \hat{p} for each possible value of \hat{p} between 0 and 1, keeping in mind that $n\hat{p}$ must be an integer.
- The distance between the midpoint of the confidence interval and one end is sometimes called the “margin of error” associated with the finite sample size. Plot the margin of error $\frac{p_+ - p_-}{2}$ as a function of \hat{p} for each possible value of \hat{p} between 0 and 1.
- Check the coverage of the confidence intervals when the true value of p is .5 by generating $N = 1,000,000$ binomial random variables each with $n = 100$ and $p = .5$. (You can do this in numpy with the command `binomial(n,p,N)`.) Determine what fraction lie below, within, and above the confidence interval (p_-, p_+) .
- Repeat part (c) for a true value of $p = .7$.